

An observation-based global monthly gridded sea surface pCO₂ product from 1998 through 2011 and its monthly climatology.

Authors:

P. Landschützer, Institute of Biogeochemistry and Pollutant Dynamics, Swiss Federal Institute of Technology (ETH), Zürich, Switzerland.

N. Gruber, Institute of Biogeochemistry and Pollutant Dynamics, Swiss Federal Institute of Technology (ETH), Zürich, Switzerland.

D.C.E. Bakker, Centre for Ocean and Atmospheric Sciences, School of Environmental Sciences, University of East Anglia, Norwich Research Park, Norwich, UK.

U. Schuster, College of Life and Environmental Sciences, Hatherly Laboratories, University of Exeter, Exeter, UK.

Citation:

This product is free to be used. Please cite the data as:

Landschützer, P., N. Gruber, D.C.E. Bakker, and U. Schuster. 2014. An observation-based global monthly gridded sea surface pCO₂ product from 1998 through 2011 and its monthly climatology. http://cdiac.ornl.gov/ftp/oceans/spco2_1998_2011_ETH_SOM-FFN. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge, Tennessee. doi: 10.3334/CDIAC/OTG.SP_CO2_1998_2011_ETH_SOM-FFN

Please cite the method as:

Landschützer, P., Gruber, N., Bakker, D. C. E., Schuster, U.: Recent variability of the global ocean carbon sink, *Global Biogeochemical Cycles*, 28, doi: 10.1002/2014GB004853, 2014

Method:

The observation-based pCO₂ fields were created using a 2-step neural network technique. In a first step, the global ocean is divided into 16 biogeochemical provinces using a self organizing map. In a second step, the non-linear relationship between variables known to drive the surface ocean carbon system and gridded observations from the SOCATv2 dataset (Bakker et al. 2014) is reconstructed using a feed-forward neural network within each province separately. The final product is then produced by projecting these driving variables, i.e., surface temperature, chlorophyll, mixed layer depth, and atmospheric CO₂ onto oceanic pCO₂ using these non-linear relationships. This results in monthly pCO₂ fields at 1°x1° resolution covering the entire globe with the exception of the Arctic Ocean and few marginal seas. The air-sea CO₂ flux is then computed using a standard bulk formula. More details can be found in Landschützer et al. 2013 and Landschützer et al. 2014.

Content:

File 1: spco2_1998-2011_ETH_SOM-FFN_CDIAC_G05.nc

The netcdf file contains:

- lat: latitude in degrees north (89.5°S – 89.5°N with 1° resolution)
- lon: longitude in degrees east (179.5°W-179.5°E with 1° resolution)
- time: time in seconds since 2000-01-01-00:00 (monthly resolution)
- spco2_raw: The raw 2-step neural network sea surface pCO₂ output as described in Landschützer et al. 2014 in μatm
- fgco2_raw: The air-sea flux density, calculated from the raw sea surface pCO₂ output as described in Landschützer et al. 2014 in $\text{mol m}^{-2} \text{yr}^{-1}$.
- spco2_smoothed: A smoothed product of the raw pCO₂, created by the spatial and temporal mean of each points neighboring pixels (the 3x3x3 pixel neighborhood domain) in μatm .
- fgco2_smoothed: A smoothed product of the raw air-sea CO₂ flux, created by the spatial and temporal mean of each points neighboring pixels (the 3x3x3 pixel neighborhood domain) in $\text{mol m}^{-2} \text{yr}^{-1}$.
- aco2: Atmospheric pCO₂ in μatm for the air-sea flux calculation, derived from the Globalview Marine Boundary Layer (MBL) xCO₂ product and SST (Reynolds et al. 2002) as well as sea level pressure (Kalnay et al. 1996) following Dickson et al. 2007.
- dco2: Delta pCO₂ calculated from the CO₂ partial pressure difference between atmosphere and the raw surface ocean partial pressure.
- dco2_smoothed: Delta pCO₂ calculated from the CO₂ partial pressure difference between atmosphere and the smoothed surface ocean partial pressure.
- kw: The gas transfer velocity calculated from the CCMP wind product (Atlas et al. 2011) as described in Landschützer et al. 2014 in m yr^{-1} .
- sol: The CO₂ solubility in $\text{mol m}^{-3} \mu\text{atm}^{-1}$ calculated from sea surface temperature (Reynolds et al. 1996) and ECCO2 sea surface salinity (Menemenlis et al. 2008) following Weiss 1974.
- ice: The percentage of sea ice from the Rayner et al. 2003 sea ice product.
- lsmask: The land-sea mask used in Landschützer et al. 2014.

File 2: spco2_clim_ETH_SOM-FFN_CDIAC_G05.nc

The netcdf file contains:

- lat: latitude in degrees north (89.5°S – 89.5°N with 1° resolution)
- lon: longitude in degrees east (179.5°W-179.5°E with 1° resolution)
- spco2_clim: The monthly pCO₂ climatology calculated from the average of the 14 year (1998-2011) smoothed neural network output in μatm .
- fgco2_clim: The monthly air-sea CO₂ flux climatology calculated from the average of the 14 year (1998-2011) smoothed neural network output in $\text{mol m}^{-2} \text{yr}^{-1}$.
- lsmask: The land-sea mask used in Landschützer et al. 2014.

Inquiries:

Inquiries should be sent to Peter Landschützer: peter.landschuetzer@usys.ethz.ch. Other data formats (e.g. matlab) available upon request.

References:

Atlas, R., Hoffman, R. N., Ardizzone, J., Leidner, S. M., Jusem, J. C., Smith, D. K., and Gombos, D.: A cross-calibrated multiplatform ocean surface wind velocity product for meteorological and oceanographic applications, *Bulletin of the American Meteorological Society*, 92, 157–174, 2011.

Bakker, D. C. E., Pfeil, B., Smith, K., Hankin, S., Olsen, A., Alin, S. R., Cosca, C., Harasawa, S., Kozyr, A., Nojiri, Y., O'Brien, K. M., Schuster, U., Telszewski, M., Tilbrook, B., Wada, C., Akl, J., Barbero, L., Bates, N. R., Boutin, J., Bozec, Y., Cai, W.-J., Castle, R. D., Chavez, F. P., Chen, L., Chierici, M., Currie, K., de Baar, H. J. W., Evans, W., Feely, R. A., Fransson, A., Gao, Z., Hales, B., Hardman-Mountford, N. J., Hoppema, M., Huang, W.-J., Hunt, C. W., Huss, B., Ichikawa, T., Johannessen, T., Jones, E. M., Jones, S. D., Jutterström, S., Kitidis, V., Körtzinger, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Manke, A. B., Mathis, J. T., Merlivat, L., Metzl, N., Murata, A., Newberger, T., Omar, A. M., Ono, T., Park, G.-H., Paterson, K., Pierrot, D., Rios, A. F., Sabine, C. L., Saito, S., Salisbury, J., Sarma, V. V. S. S., Schlitzer, R., Sieger, R., Skjelvan, I., Steinhoff, T., Sullivan, K. F., Sun, H., Sutton, A. J., Suzuki, T., Sweeney, C., Takahashi, T., Tjiputra, J., Tsurushima, N., van Heuven, S. M. A. C., Vandemark, D., Vlahos, P., Wallace, D. W. R., Wanninkhof, R., and Watson, A. J.: An update to the Surface Ocean CO₂ Atlas (SOCAT version 2), *Earth System Science Data*, 6, 69–90, doi:10.5194/essd-6-69-2014, URL: <http://www.earth-syst-sci-data.net/6/69/2014/>, 2014.

Dickson, A. G., Sabine, C. L., and Christian, J. R., eds.: *Guide to Best Practices for Ocean CO₂ Measurements*, PICES Special Publication, IOCCP Report No. 8, 2007.

GLOBALVIEW-CO₂: Cooperative Atmospheric Data Integration Project - Carbon Dioxide, CD-ROM, NOAA ESRL, Boulder, Colorado [Also available on Internet via anonymous FTP to <ftp.cmdl.noaa.gov>, Path: ccg/co2/GLOBALVIEW], 2014.

Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Leetmaa, A., Reynolds, R., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Jenne, R., and Joseph, D.: The NCEP/NCAR 40-year reanalysis project, *Bulletin of the American Meteorological Society*, 77, 437–470, 1996.

Landschützer, P., Gruber, N., Bakker, D. C. E., Schuster, U.: Recent variability of the global ocean carbon sink, *Global Biogeochemical Cycles*, 28, doi: 10.1002/2014GB004853, 2014.

Landschützer, P., Gruber, N., Bakker, D. C. E., Schuster, U., Nakaoka, S., Payne, M. R., Sasse, T., and Zeng, J.: A neural network-based estimate of the seasonal to inter-annual variability of the Atlantic Ocean carbon sink, *Biogeosciences*, 10, 7793–7815, doi:10.5194/bg-10-7793-2013, 2013.

Menemenlis, D., Campin, J., Heimbach, P., Hill, C., Lee, T., Nguyen, A., Schodlok, M., and Zhang, H.: ECCO2: High resolution global ocean and sea ice data synthesis, *Mercator Ocean Quarterly Newsletter*, 31, 13–21, 2008.

Rayner, N. A., Parker, D. E., Horton, E. B., Folland, C. K., Alexander, L. V., Rowell, D. P., Kent, E. C., and Kaplan, A.: Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century, *Journal of Geophysical Research*, 108, 4407, doi:10.1029/2002JD002670, 2003.

Reynolds, R. W., Rayner, N. A., Smith, T. M., Stokes, D. C., and Wang, W.: An improved in situ and satellite SST analysis for climate, *Journal of Climate*, 15, 1609–1625, 2002.

Weiss, R. F.: Carbon Dioxide in Water and Seawater: The Solubility of a Non-Ideal Gas, *Marine Chemistry*, 2, 203–215, 1974.