

## **XBT Data Processing – PODS 1992 DSJ**

XBT data were collected with SEAS software (Shipboard Environmental data Acquisition System, NOAA/AOML) on the NOAA Ship David Starr Jordan, Aug 8 through Oct 25, 1992. Typically, XBT drops were conducted every three hours during daylight operations. Several types of Sippican Expendable Bathythermograph (XBT) probes were used on this cruise to measure the temperature of the water column. Type T-7 measures to 760 m, T-4 and T-6 measure to 460 m, and T-10 measures to 200 m.

Raw data files were checked for date/time and position and then edited to remove erroneous temperature data. All valid and edited profiles were then combined into final files. The XBTCorrect file has depth and temperature corrections according to Cheng et al. (2016, Bull. American Meteorological Soc., June 2016, 923-933). The XBTArchive file has depths calculated with the fall rate equation of Hanawa et al. (1995, Deep-Sea Research I 42:1423-1451) and excludes incomplete profiles that do not extend through the thermocline. This file is for archiving and allows for optional correction of depth and temperature by users.

All data and programs are stored in sub-folders of P:\Surveys\Data\ETP\1992\_PODS\Oceanographic\XBT\.

The raw data files are in the sub-folder Raw. The programs used to process the data are stored in the sub-folder Current\_Version\Programs. These programs read year-specific information from \*.ini files, which are stored in the same folder as the programs. The intermediate files created during the processing steps are stored in the sub-folder Current\_Version\Intermediate. The final processed data and documentation are stored in the sub-folder Current\_Version\Final.

### **Raw Data Files**

SEAS raw data files were converted to \*.edf files with Fortran program SEAS2EDF. No \*.ini file is used by the SEAS2EDF program.

The name of the raw SEAS files start with a 2-digit code that appears to tell the type of probe that was used: 04 = T-4, 06 = T-6, 07 = T-7, and 10 = T-10. This is consistent with the depths that those probes went to. Therefore, the Probe Type was set based on the code in the file name.

### **Raw File Summaries**

The data collected on each survey were backed up on several DVD's. The DVDs were frequently created at the end of each leg. Sometimes data from each leg were backed up separately, while at other times the backups included all data collected (e.g., data collected on legs 1-3 were backed up at the end of leg 3) . Consequently, there can be a great deal of overlap among the DVDs.

However, for PODS 1992, there were no DVD's. The data was found in a folder on the network: W:\Final\_Oceanographic\ETP\1992\_PODS\Oceanographic\XBT\Raw\DSJ. These files were copied to P:\Surveys\Data\ETP\1992\_PODS\Oceanographic\XBT\Raw\DSJ\.

No XBT spreadsheets or Elog databases were created during the cruise.

Perl program XBTRawFileInfo was used to get a summary of the raw files. It showed that all SEAS raw files had matching EDF Files, and vice versa.

## **Data Checks**

### Position Check

Perl program XBTPositionCheck compares the positions and times of the XBT drops to positions and times in the final TSG "Track Check" file. The program reads the following information from its .ini file: the location of the EDF files, the location of the final TSG Track Check file, and the number of seconds to search the TSG file for a record that matches each XBT file's time. Positions for stations without corresponding TSG data are checked against the XBT Log and ODL.

For 1992 PODS DSJ, the most glaring problem was that Longitude was positive, which is incorrect for the Western hemisphere. The Longitudes were manually negated directly in the EDF files. Once that was fixed, XBTPositionCheck was run again. Seven drops' positions were greater than 1 nautical mile from TSG positions matched by time. For five of these, the XBT position appeared to be incorrect. These were corrected directly in the EDF files. For the remaining two, it was impossible to tell which was correct. Since they were only slightly over 1 nautical mile from TSG, they were accepted.

### Station and Drop Lists

Perl program XBTPositionAndDrops creates the Station and Drops files. From its ini file, it reads the location of the EDF files, and the location of the time offsets file. The Drop file is simply a list of the EDF Files. The Station file lists additional information for each drop: time and location, number of data points, probe type, start and end depths, and drop rate coefficients in the EDF file.

For 1992 PODS DSJ, there were 135 drops. 78 used T-4 probes, 22 used T-6 probes, 34 used T-7 probes, and 1 used a T-10 probe.

### Profile Review and Edit

At the lab, profiles were examined by Paul Fiedler using the Visual Basic program ProfileViewer, which is located in

P:\Surveys\Data\_Processing\_Resources\Oceanographic\XBT\ProfileViewer. The reviewer decides what edits need to be made, and records that information in file XBTDrops\_1992\_PODS\_DSJ.txt. Each XBT drop has a line in that file, with the EDF file name. If the profile requires any editing, the reviewer added codes for the edits to be applied. Codes

are based on Bailey et al. (Quality Control Cookbook for XBT Data, CSIRO Marine Laboratories Report 221, 1994). The Quality Control Cookbook can be found in P:\Surveys\Data\_Processing\_Resources\Oceanographic\XBT\ . Codes are as follows, where ddd.d is the depth exactly as recorded in the .edf file:

- >RJ                      Reject profile.
  
- >NN                      The profile is not suitable for submission to the NCEI World Ocean Database, but is good enough for deriving surface temperature and mixed layer depth, and perhaps thermocline variables, for MMTD purposes.
  
- >XD ddd.d ddd.d        Exclude data within the profile. Reject data between the first and second specified depths.
  
- >BD ddd.d                Exclude data at the bottom end of the profile. Reject data deeper than ddd.d.
  
- >FT ddd.d                False trigger, data recorded before the probe entered the water. This error is rare. Reject all data above ddd.d and change the depth at ddd.d to 0.7m, which is the starting depth for all drops (and then increment the depth estimates per Hanawa et al. 1995).
  
- >ST ddd.d                Surface transient, identified by warming or cooling as the probe temperature equilibrates. Reject data before depth ddd.d. All temperatures at depths  $\leq 4.0$  m are rejected to routinely dismiss surface transients, so this flag is only effective if a depth  $>4.0$  m is indicated. If the profile also contains an FT error, ddd.d is an integer that indicates the number of records after the FT correction that should be rejected.

Profile rejections and data exclusions are usually explained in a comment following an apostrophe ('): wb = wire break, ws = wire stretch, hb = the probe hit the bottom, ip = insulation penetration.

### Apply Profile Edits

Perl program XBTEdit reads the editing codes entered by the reviewer, and creates edited output files. From its ini file, it reads: The location of the edit file (created in the previous step), the location of the time offsets file, the location of the raw XBT files, and the coefficients for the fall rate equation. It creates output files with the editing codes applied. (e.g. Drops marked with RJ are excluded, depths marked with XD are removed, etc.)

Here is the output from XBTEdit:

```
135        lines read from input file
          P:\Surveys\Data\ETP\1992_PODS\Oceanographic\XBT\Current_Version\Intermed
          iate\XBTDrops_1992_PODS_DSJ.txt
```

```

0      lines started with an apostrophe, so were ignored
14     lines had an RJ code, so were ignored

121    records written to
       P:\Surveys\Data\ETP\1992_PODS\Oceanographic\XBT\Current_Version\Intermed
       iate\XBTEdit_1992_PODS_DSJ.dat

3      lines were not good enough for NODC, so their drop numbers were negated
1      records written to
       P:\Surveys\Data\ETP\1992_PODS\Oceanographic\XBT\Current_Version\Intermed
       iate\XBTNODC_1992_PODS_DSJ_6.301_0.00216.dat
117    records written to
       P:\Surveys\Data\ETP\1992_PODS\Oceanographic\XBT\Current_Version\Intermed
       iate\XBTNODC_1992_PODS_DSJ_6.691_0.00225.dat

12.30  Average temperature of all 93028 output points whose drops met the Min
       To Avg depth
20.95  Average temperature of all 17264 output points <= 100 m whose drops met
       the Min To Avg depth
       Note that this is based on depths calculated using the Hanawa et al 1995
       formula

```

## Depth and Temperature Corrections

Perl program XBTCorrect was used to correct depths and temperatures, using the method described in P:\Surveys\Data\_Processing\_Resources\Oceanographic\XBT\CH XBT fall-rate method.pdf. Here is the output from XBTCorrect:

```

Reading XBTCorrect.ini...
Reading CH14 table 'P:\Surveys\Data_Processing_Resources\Oceanographic\XBT\Fall
rate\CH14_table1.txt'...
Reading CH14 table 'P:\Surveys\Data_Processing_Resources\Oceanographic\XBT\Fall
rate\CH14_table2.txt'...
Processing
       P:\Surveys\Data\ETP\1992_PODS\Oceanographic\XBT\Current_Version\Intermed
       iate\XBTEdit_1992_PODS_DSJ.dat...
T-4: Thermal Bias = 0.05864, A = 6.786, B = 0.00332, Offset = 2.015
T-7: Thermal Bias = 0.01222, A = 6.661, B = 0.00263, Offset = 2.181
T-6: Thermal Bias = 0.05864, A = 6.786, B = 0.00332, Offset = 2.015
T-10: Thermal Bias = 0.04244, A = 6.503, B = 0.00137, Offset = 0.376

122    lines read from
       'P:\Surveys\Data\ETP\1992_PODS\Oceanographic\XBT\Current_Version\Interme
       diate\XBTEdit_1992_PODS_DSJ.dat'
0      lines started with an apostrophe, so were ignored
31     records written to
       P:\Surveys\Data\ETP\1992_PODS\Oceanographic\XBT\Current_Version\Intermed
       iate\XBTCorrect_1992_PODS_DSJ_T-7.dat
1      records written to
       P:\Surveys\Data\ETP\1992_PODS\Oceanographic\XBT\Current_Version\Intermed
       iate\XBTCorrect_1992_PODS_DSJ_T-10.dat
69     records written to
       P:\Surveys\Data\ETP\1992_PODS\Oceanographic\XBT\Current_Version\Intermed
       iate\XBTCorrect_1992_PODS_DSJ_T-4.dat
20     records written to
       P:\Surveys\Data\ETP\1992_PODS\Oceanographic\XBT\Current_Version\Intermed
       iate\XBTCorrect_1992_PODS_DSJ_T-6.dat

```

3        had negative drop numbers, which means their data was not good enough  
         for NODC

## Final data files

The NODC files created in the XBTEdit step were copied to \Final and were renamed as:

XBTArchive\_1992\_PODS\_DSJ\_T-4\_T-6\_T-7.dat  
XBTArchive\_1992\_PODS\_DSJ\_T-10.dat

Each drop record in a XBTArchive file starts with fields under the following column headers:

UTC Date, UTC Time, Time Offset, Local Date, Local Time, Latitude, Longitude,  
Drop #, Probe Type, # Points, H95 FRE Coefficients...

The column headers after that are depths (m), based on the Hanawa et al 1995 fall rate equation,  
e.g:

..., 0.7, 1.3, 2.0, 2.7, 3.3, 4.0, 4.7, 5.4, 6.0, 6.7, 7.4, 8.0, 8.7, ...

The drop records data in those columns are temperatures (°C) at those depths. Missing data (e.g.,  
surface transients) are blank.

The files that had depth and temperature corrected based on Cheng et al were also copied to  
\Final. Those files are:

XBTCorrect\_1992\_PODS\_DSJ\_T-4.dat  
XBTCorrect\_1992\_PODS\_DSJ\_T-6.dat  
XBTCorrect\_1992\_PODS\_DSJ\_T-7.dat  
XBTCorrect\_1992\_PODS\_DSJ\_T-10.dat

Each drop record in a XBTCorrect file starts with fields under the following column headers:

UTC Date, UTC Time, Time Offset, Local Date, Local Time, Latitude, Longitude,  
Drop #, Probe Type, CH14 Values, # Points...

The column headers after that are depths (m), based on the Cheng et al fall rate correction, e.g:

..., 0.5, 1.1, 1.8, 2.5, 3.1, 3.8, 4.5, 5.1, 5.8, 6.5, 7.1, 7.8, 8.5, 9.1, ...

Paul Fiedler  
Dan Prosper  
01 June 2021