XBT Data Processing – HICEAS 2002 MAC

XBT data were collected with SEAS software (Shipboard Environmental data Acquisition System, NOAA/AOML) on the NOAA Ship McArthur, Oct 11 through Dec 9, 2002. Typically, XBT drops were conducted daily at 0900, 1200, 1500 hours (local ship time), and after sunset. Sippican Deep Blue Expendable Bathythermograph (XBT) probes were used to measure the temperature of the water column to 760 m.

Raw data files were checked for date/time and position and then edited to remove erroneous temperature data. All valid and edited profiles were then combined into final files. The XBTCorrect file has depth and temperature corrections according to Cheng et al. (2016, Bull. American Meteorological Soc., June 2016, 923-933). The XBTArchive file has depths calculated with the fall rate equation of Hanawa et al. (1995, Deep-Sea Research I 42:1423-1451) and excludes incomplete profiles that do not extend through the thermocline. This file is for archiving and allows for optional correction of depth and temperature by users.

All data and programs are stored in sub-folders of

P:\Surveys\Data\PacificIslands\2002_HICEAS\Oceanographic\XBT\. The raw data files are in the sub-folder Raw. The programs used to process the data are stored in the sub-folder Current_Version\Programs. These programs read year-specific information from *.ini files, which are stored in the same folder as the programs. The intermediate files created during the processing steps are stored in the sub-folder Current_Version\Intermediate. The final processed data and documentation are stored in the sub-folder Current_Version\Final.

Raw Data Files

SEAS raw data files were converted to *.edf files with Fortran program SEAS2EDF. No *.ini file is used by the SEAS2EDF program.

Raw File Summaries

The data collected on each survey were backed up on several DVD's. The DVDs were frequently created at the end of each leg. Sometimes data from each leg were backed up separately, while at other times the backups included all data collected (e.g., data collected on legs 1-3 were backed up at the end of leg 3). Consequently, there can be a great deal of overlap among the DVDs. For the data collected on the MAC during HICEAS 2002, the CD labeled HICEAS2002_Legs12_McArthur was the most complete. However, not all files were on that DVD. These files were copied to P:\Surveys\Data\ETP\2003 STAR\Oceanographic\XBT\Raw\MACII\.

Information about XBT drops, such as time, date, latitude, longitude, and comments about the profile by the at-sea oceanographer, was recorded at sea in an Excel spreadsheet called XBT02 processing.xls. However, this was only done for drops on 10/28/2002 and later.

Perl program XBTRawFileInfo was used to get a summary of the raw files. It showed that the drop numbers were re-set to 1 several times during the cruise. Therefore, the EDF files were manually edited to match the drop numbers in file XBT02_processing.xls. This also gives all drops consecutive numbers for the entire cruise.

- In Leg 1, the first set of drops were numbered 1 to 68. These were left unchanged
- In Leg 1, the second set of drops 1 through 33 were increased by 68, changing them to 69 to 101
- In Leg 2, 101 was added to all drop numbers (1 to 61), changing them to 102 to 162
- In Leg 3, 162 was added to all drop numbers (1 to 47), changing them to 163 to 209

Note that there was another problem with drop numbers in the ODL. There are two drop files from 10/28 around 19:30. Thos drops' files were originally numbered 001 and 002. They were re-numbered above to be 069 and 070. However, there is only one drop in the ODL around that time, and it's numbered "069 / 002". For the remainder of Leg 1, the cruise-wide drop numbers in the ODL (69 through 100) are one lower than the numbers they were given above (69 to 101). When looking for a drop in the ODL, subtract one from the number of the file.

Data Checks

Position Check

Perl program XBTPositionCheck compares the positions and times of the XBT drops to positions and times in the final TSG "Track Check" file. The program reads the following information from its .ini file: the location of the EDF files, the location of the final TSG Track Check file, and the number of seconds to search the TSG file for a record that matches each XBT file's time. Positions for stations without corresponding TSG data are checked against the XBT Log and ODL.

```
209
               EDF files read
               were missing date, time, latitude, or longitude
          0
          0
               were not in a valid format
          8
             did not have a TSG record within 120 seconds in
P:\Surveys\Data\PacificIslands\2002 HICEAS\Oceanographic\TSG\Current Version\In
termediate\TSGTrackCheck 2002 HICEAS MAC Final.dat
         201
               lines were compared to TSG
    0.47510
             Root-mean-square-error of distances
               Distances >= 1 nautical mile
         63
     3.58630
               Max distance
```

For 200 HICEAS MAC, 63 XBT drops were greater than 1 nautical mile from TSG positions matched by time. Those whose position and time in the ODL more closely matched TSG were corrected based on the ODL. However, the ODL did not have positions for some drops, so not all could be corrected.

Station and Drop Lists

Perl program XBTStationAndDrops creates the Station and Drops files. From its ini file, it reads the location of the EDF files, and the location of the time offsets file. The Drop file is simply a list of the EDF Files. The Station file lists additional information for each drop: time and location, number of data points, probe type, start and end depths, and drop rate coefficients in the EDF file.

For 2002 HICEAS MAC, there were 209 drops, all using Deep Blue probes.

Profile Review and Edit

At the lab, profiles were examined by Paul Fiedler using the Visual Basic program ProfileViewer, which is located in P:\Surveys\Data_Processing_Resources\Oceanographic\XBT\ProfileViewer. The reviewer decides what edits need to be made, and records that information in file XBTDrops_2002_HICEAS_MAC.txt. Each XBT drop has a line in that file, with the EDF file name. If the profile requires any editing, the reviewer added codes for the edits to be applied. Codes are based on Bailey et al. (Quality Control Cookbook for XBT Data, CSIRO Marine Laboratories Report 221, 1994). The Quality Control Cookbook can be found in P:\Surveys\Data_Processing_Resources\Oceanographic\XBT\. Codes are as follows, where ddd.d is the depth exactly as recorded in the .edf file:

| >RJ | Reject profile. |
|-----------------|--|
| >NN | The profile is not suitable for submission to the NCEI World Ocean Database, but is good enough for deriving surface temperature and mixed layer depth, and perhaps thermocline variables, for MMTD purposes. |
| >XD ddd.d ddd.d | Exclude data within the profile. Reject data between the first and second specified depths. |
| >BD ddd.d | Exclude data at the bottom end of the profile. Reject data deeper than ddd.d. |
| >FT ddd.d | False trigger, data recorded before the probe entered the water. This error is rare. Reject all data above ddd.d and change the depth at ddd.d to 0.7m, which is the starting depth for all drops (and then increment the depth estimates per Hanawa et al. 1995). |
| >ST ddd.d | Surface transient, identified by warming or cooling as the probe temperature equilibrates. Reject data before depth ddd.d. All temperatures at depths \leq 4.0 m are rejected to routinely dismiss surface transients, so this flag is only effective if a depth >4.0 m is indicated. If the profile also contains an FT error, ddd.d is an integer that indicates the number of |

records after the FT correction that should be rejected.

Profile rejections and data exclusions are usually explained in a comment following an apostrophe ('): wb = wire break, ws = wire stretch, hb = the probe hit the bottom, ip = insulation penetration.

Apply Profile Edits

Perl program XBTEdit reads the editing codes entered by the reviewer, and creates edited output files. From its ini file, it reads: The location of the edit file (created in the previous step), the location of the time offsets file, the location of the raw XBT files, and the coefficients for the fall rate equation. It creates output files with the editing codes applied. (e.g. Drops marked with RJ are excluded, depths marked with XD are removed, etc.)

The program reported the following:

```
209lines read from input file
P:\Surveys\Data\PacificIslands\2002_HICEAS\Oceanographic\XBT\Current_Version
\Intermediate\XBTDrops_2002_HICEAS_MAC.txt
0 lines started with an apostrophe, so were ignored
6 lines had an RJ code, so were ignored
203records written to
P:\Surveys\Data\PacificIslands\2002_HICEAS\Oceanographic\XBT\Current_Version
\Intermediate\XBTEdit_2002_HICEAS_MAC.dat
3 lines were not good enough for NODC, so their drop numbers were negated
200records written to
P:\Surveys\Data\PacificIslands\2002_HICEAS\Oceanographic\XBT\Current_Version
\Intermediate\XBTEdit_2002_HICEAS_MAC.dat
3 lines were not good enough for NODC, so their drop numbers were negated
200records written to
P:\Surveys\Data\PacificIslands\2002_HICEAS\Oceanographic\XBT\Current_Version
\Intermediate\XBTNODC_2002_HICEAS_MAC_6.691_0.00225.dat
11.88 Average temperature of all 221237 output points whose drops met the Min
To Avg depth
```

```
24.30 Average temperature of all 29356 output points <= 100 m whose drops met
the Min To Avg depth
Note that this is based on depths calculated using the Hanawa et al 1995
formula
```

Correct Depth and Temperature

Perl program XBTCorrect was used to correct depths and temperatures, using the method described in P:\Surveys\Data_Processing_Resources\Oceanographic\XBT\CH XBT fall-rate method.pdf. The program reported the following:

```
Reading XBTCorrect.ini...
Reading CH14 table 'P:\Surveys\Data_Processing_Resources\Oceanographic\XBT\Fall
rate\CH14_table1.txt'...
Reading CH14 table 'P:\Surveys\Data_Processing_Resources\Oceanographic\XBT\Fall
rate\CH14 table2.txt'...
```

```
Processing
P:\Surveys\Data\PacificIslands\2002_HICEAS\Oceanographic\XBT\Current_Version
\Intermediate\XBTEdit_2002_HICEAS_MAC.dat...
Deep Blue: Thermal Bias = 0.05143, A = 6.622, B = 0.00235, Offset = 1.932
204 lines read from
'P:\Surveys\Data\PacificIslands\2002_HICEAS\Oceanographic\XBT\Current_Versio
n\Intermediate\XBTEdit_2002_HICEAS_MAC.dat'
0 lines started with an apostrophe, so were ignored
203 records written to
P:\Surveys\Data\PacificIslands\2002_HICEAS\Oceanographic\XBT\Current_Version
\Intermediate\XBTCorrect_2002_HICEAS_MAC_Deep Blue.dat
3 had negative drop numbers, which means their data was not good enough for
NODC
```

Final data files

The NODC file created in the XBTEdit step was copied to \Final and was renamed as: XBTArchive 2002 HICEAS MAC Deep Blue.dat

Each drop record in a XBTArchive file starts with fields under the following column headers:

UTC Date, UTC Time, Time Offset, Local Date, Local Time, Latitude, Longitude, Drop #, Probe Type, # Points, H95 FRE Coefficients...

The column headers after that are depths (m), based on the Hanawa et al 1995 fall rate equation, e.g:

..., 0.7, 1.3, 2.0, 2.7, 3.3, 4.0, 4.7, 5.4, 6.0, 6.7, 7.4, 8.0, 8.7,...

The drop records data in those columns are temperatures (°C) at those depths. Missing data (e.g., surface transients) are blank.

The file that had depth and temperature corrected based on Cheng et al was also copied to \Final. This file is:

XBTCorrect 2002 HICEAS MAC Deep Blue.dat

Each drop record in a XBTCorrect file starts with fields under the following column headers:

UTC Date, UTC Time, Time Offset, Local Date, Local Time, Latitude, Longitude, Drop #, Probe Type, CH14 Values, # Points...

The column headers after that are depths (m), based on the Cheng et al fall rate correction, e.g.

..., 0.1, 0.7, 1.4, 2.0, 2.7, 3.4, 4.0, 4.7, 5.3, 6.0, 6.7, 7.3, 8.0, 8.7,...

Paul Fiedler Dan Prosperi

01 June 2021