

## **XBT Data Processing – PICEAS 2005 MACII**

XBT data were collected with SEAS software (Shipboard Environmental data Acquisition System, NOAA/AOML) on the NOAA Ship McArthur II, 7/29/2015 through 11/29/2015. Typically, XBT drops were conducted every three hours during daylight operations at 0900, 1200, and 1500 (local ship time). Some XBT drops were performed in lieu of CTD casts at 0400 or 2000 (local ship time). Sippican Deep Blue Expendable Bathythermograph (XBT) probes measure the temperature of the water column to 760 m.

Raw data files were checked for date/time and position and then edited to remove erroneous temperature data. All valid and edited profiles were then combined into final files. The XBTCorrect file has depth and temperature corrections according to Cheng et al. (2016, Bull. American Meteorological Soc., June 2016, 923-933). The XBTArchive file has depths calculated with the fall rate equation of Hanawa et al. (1995, Deep-Sea Research I 42:1423-1451) and excludes incomplete profiles that do not extend through the thermocline. This file is for archiving and allows for optional correction of depth and temperature by users.

All data and programs are stored in sub-folders of P:\Surveys\Data\PacificIslands\2005\_PICEAS\Oceanographic\XBT\ . The raw data files are in the sub-folder Raw. The programs used to process the data are stored in the sub-folder Current\_Version\Programs. These programs read year-specific information from \*.ini files, which are stored in the same folder as the programs. The intermediate files created during the processing steps are stored in the sub-folder Current\_Version\Intermediate. The final processed data and documentation are stored in the sub-folder Current\_Version\Final.

### **Raw Data Files**

SEAS raw data files were converted to \*.edf files with Fortran program SEAS2EDF. No \*.ini file is used by the SEAS2EDF program.

### **Raw File Summaries**

The data collected on each survey were backed up on several DVD's. The DVDs were frequently created at the end of each leg. Sometimes data from each leg were backed up separately, while at other times the backups included all data collected (e.g., data collected on legs 1-3 were backed up at the end of leg 3) . Consequently, there can be a great deal of overlap among the DVDs.

For the data collected on the MACII during PICEAS 2005, the CD labeled CD\_PICEAS\_2005 was the most complete. These files were identical to files already on P:\Surveys\Data\PacificIslands\2005\_PICEAS\Oceanographic\XBT\Raw\MACII\.

No XBT spreadsheets or Elog databases were created during the cruise.

Perl program XBTRawFileInfo was used to get a summary of the raw files. It showed that all SEAS files had matching EDF Files, and vice versa.

## Data Checks

### Position Check

Perl program XBTPositionCheck compares the positions and times of the XBT drops to positions and times in the final TSG "Track Check" file. The program reads the following information from its .ini file: the location of the EDF files, the location of the final TSG Track Check file, and the number of seconds to search the TSG file for a record that matches each XBT file's time. Positions for stations without corresponding TSG data are checked against the XBT Log and ODL.

For 2005 PICEAS, many XBT drops were greater than 1 nautical mile from TSG positions matched by time. The positions of 12 of these were corrected in the EDF files based on the position written in the ODL, or the TSG position. These RDF files were moved to the Files\_With\_Errors sub-folder of the Raw folder. The remaining position differences were apparently because of time differences between XBT and TSG. The time in five files was corrected based on the time in the ODL. Drop 190 was 2.7 nautical miles from TSG based on time, but the time and position matched the ODL, so it was accepted. The remaining were all within 2.5 nautical miles, and were all accepted.

### Results after correcting the above:

```
354      EDF files read
      0      were missing date, time, latitude, or longitude
      0      were not in a valid format
      0      did not have a TSG record within 120 seconds in
P:\Surveys\Data\PacificIslands\2005_PICEAS\Oceanographic\TSG\Current_Version\In
termediate\MACII\TSGTrackCheck_2005_PICEAS_MACII.dat

354      lines were compared to TSG

0.54461      Root-mean-square-error of distances
139      Distances >= 1 nautical mile
3.61390      Max distance
```

### Station and Drop Lists

Perl program XBTPositionAndDrops creates the Station and Drops files. From its ini file, it reads the location of the EDF files, and the location of the time offsets file. The Drop file is simply a list of the EDF Files. The Station file lists additional information for each drop: time and location, number of data points, probe type, start and end depths, and drop rate coefficients in the EDF file.

For 2005 PICEAS, there were 354 valid drops, all using Deep Blue probes.

### Profile Review and Edit

At the lab, profiles were examined by Paul Fiedler using the Visual Basic program ProfileViewer, which is located in

P:\Surveys\Data\_Processing\_Resources\Oceanographic\XBT\ProfileViewer. The reviewer decided what edits needed to be made, and recorded that information in file XBTedit05m.txt. Each XBT drop has a line in that file, with the EDF file name. If the profile requires any editing, the reviewer added codes for the edits to be applied. Codes are based on Bailey et al. (Quality Control Cookbook for XBT Data, CSIRO Marine Laboratories Report 221, 1994). The Quality Control Cookbook can be found in

P:\Surveys\Data\_Processing\_Resources\Oceanographic\XBT\ . Codes are as follows, where ddd.d is the depth exactly as recorded in the .EDF file:

- |                 |   |
|-----------------|---|
| >RJ             | Reject profile.   |
| >NN             | The profile is not suitable for submission to the NCEI World Ocean Database, but is good enough for deriving surface temperature and mixed layer depth, and perhaps thermocline variables, for MMTD purposes.   |
| >XD ddd.d ddd.d | Exclude data within the profile. Reject data between the first and second specified depths.   |
| >BD ddd.d       | Exclude data at the bottom end of the profile. Reject data deeper than ddd.d.   |
| >FT ddd.d       | False trigger, data recorded before the probe entered the water. This error is rare. Reject all data above ddd.d and change the depth at ddd.d to 0.7m, which is the starting depth for all drops (and then increment the depth estimates per Hanawa et al. 1995).  |
| >ST ddd.d       | Surface transient, identified by warming or cooling as the probe temperature equilibrates. Reject data before depth ddd.d. All temperatures at depths $\leq 4.0$ m are rejected to routinely dismiss surface transients, so this flag is only effective if a depth $>4.0$ m is indicated. If the profile also contains an FT error, ddd.d is an integer that indicates the number of records after the FT correction that should be rejected. |

Profile rejections and data exclusions are usually explained in a comment following an apostrophe ('): wb = wire break, ws = wire stretch, hb = the probe hit the bottom, ip = insulation penetration.

## Apply Profile Edits

Perl program XBTEdit reads the editing codes entered by the reviewer, and creates edited output files. From its ini file, it reads: The location of the edit file (created in the previous step), the location of the time offsets file, the location of the raw XBT files, and the coefficients for the fall rate equation. It creates output files with the editing codes applied. (e.g. Drops marked with RJ are excluded, depths marked with XD are removed, etc.) Results:

```
354   lines read from input file
      P:\Surveys\Data\PacificIslands\2005_PICEAS\Oceanographic\XBT\Current_Version\Intermediate\XBTDrops_2005_PICEAS_MACII PF.txt
0     lines started with an apostrophe, so were ignored
22    lines had an RJ code, so were ignored

332   records written to
      P:\Surveys\Data\PacificIslands\2005_PICEAS\Oceanographic\XBT\Current_Version\Intermediate\XBTEdited_2005_PICEAS_MACII.dat

11    lines were not good enough for NODC, so their drop numbers were negated
321   records written to
      P:\Surveys\Data\PacificIslands\2005_PICEAS\Oceanographic\XBT\Current_Version\Intermediate\XBTNODC_2005_PICEAS_MACII_6.691_0.00225.dat

11.78 Average temperature of all 361093 output points whose drops met the Min To Avg depth
25.74 Average temperature of all 47935 output points <= 100 m whose drops met the Min To Avg depth
      Note that this is based on depths calculated using the Hanawa et al 1995 formula
```

## Correct Depth and Temperature

XBT depths and temperatures were adjusted using program XBTCorrect. This program adjusts depths and temperatures based on the method described in P:\Surveys\Data\_Processing\_Resources\Oceanographic\XBT\CH XBT fall-rate method.pdf.

Results:

```
Reading XBTCorrect.ini...
Reading CH14 table 'P:\Surveys\Data_Processing_Resources\Oceanographic\XBT\Fall rate\CH14_table1.txt'...
Reading CH14 table 'P:\Surveys\Data_Processing_Resources\Oceanographic\XBT\Fall rate\CH14_table2.txt'...
Processing
      P:\Surveys\Data\PacificIslands\2005_PICEAS\Oceanographic\XBT\Current_Version\Intermediate\XBTEdited_2005_PICEAS_MACII.dat...
Deep Blue: Thermal Bias = 0.05699, A = 6.625, B = 0.00238, Offset = 1.951

333   lines read from
      'P:\Surveys\Data\PacificIslands\2005_PICEAS\Oceanographic\XBT\Current_Version\Intermediate\XBTEdited_2005_PICEAS_MACII.dat'
0     lines started with an apostrophe, so were ignored
```

332 records written to  
P:\Surveys\Data\PacificIslands\2005\_PICEAS\Oceanographic\XBT\Current\_Version\Intermediate\XBTCorrect\_2005\_PICEAS\_MACII\_Deep Blue.dat

11 had negative drop numbers, which means their data was not good enough for NODC

## Final data files

The NODC file created in the XBTEdit step was copied to \Final and was renamed as:  
XBTArchive\_2005\_PICEAS\_MACII\_Deep Blue.dat

Each drop record in a XBTArchive file starts with fields under the following column headers:

UTC Date, UTC Time, Time Offset, Local Date, Local Time, Latitude, Longitude,  
Drop #, Probe Type, # Points, H95 FRE Coefficients...

The column headers after that are depths (m), based on the Hanawa et al 1995 fall rate equation, e.g:

..., 0.7, 1.3, 2.0, 2.7, 3.3, 4.0, 4.7, 5.4, 6.0, 6.7, 7.4, 8.0, 8.7,...

The drop records data in those columns are temperatures (°C) at those depths. Missing data (e.g., surface transients) are blank.

The file that had depth and temperature corrected based on Cheng et al was also copied to \Final. This file is:

XBTCorrect\_2005\_PICEAS\_MACII\_Deep Blue.dat

Each drop record in a XBTCorrect file starts with fields under the following column headers:

UTC Date, UTC Time, Time Offset, Local Date, Local Time, Latitude, Longitude,  
Drop #, Probe Type, CH14 Values, # Points...

The column headers after that are depths (m), based on the Cheng et al fall rate correction, e.g:

..., 0.0, 0.7, 1.4, 2.0, 2.7, 3.3, 4.0, 4.7, 5.3, 6.0, 6.7, 7.3, 8.0, 8.6,...

Paul Fiedler  
Dan Prospero  
24 May 2021