

XBT Data Processing – HICEAS 2010 MACII

XBT data were collected with Sippican MK21 software on the NOAA Ship McArthur II, 4 August through 10 December 2010. Typically, XBT drops were conducted every three hours during daylight operations at 0900, 1200, and 1500 (local ship time). Some XBT drops were performed in lieu of CTD casts at 0400 or 2000 (local ship time). Sippican Deep Blue Expendable Bathythermograph (XBT) probes measured the temperature of the water column to 760 m.

Raw data files were checked for date/time and position and then edited to remove erroneous temperature data. All valid and edited profiles were then combined into final files. The XBTCorrect file has depth and temperature corrections according to Cheng et al. (2016, Bull. American Meteorological Soc., June 2016, 923-933). The XBTArchive file has depths calculated with the fall rate equation of Hanawa et al. (1995, Deep-Sea Research I 42:1423-1451) and excludes incomplete profiles that do not extend through the thermocline. This file is for archiving and allows for optional correction of depth and temperature by users.

All data and programs are stored in sub-folders of P:\Surveys\Data\PacificIslands\2010_HICEAS\Oceanographic\XBT\ . The raw data files are in the sub-folder Raw\MACII. The programs used to process the data are stored in the sub-folder Current_Version\Programs. These programs read year-specific information from *.ini files, which are stored in the same folder as the programs. The intermediate files created during the processing steps are stored in the sub-folder Current_Version\Intermediate\MACII. The final processed data and documentation are stored in the sub-folder Current_Version\Final.

Raw Data Files

Sippican raw (*.RDF) and export (*.EDF) data files are in P:\Surveys\Data\PacificIslands\2010_HICEAS\Oceanographic\XBT\Raw\MACII.

Raw File Summaries

The data collected on each survey were backed up on several DVD's. The DVDs were frequently created at the end of each leg. Sometimes data from each leg were backed up separately, while at other times the backups included all data collected (e.g., data collected on legs 1-3 were backed up at the end of leg 3) . Consequently, there can be a great deal of overlap among the DVDs. For the data collected on the MACII during HICEAS 2010, the Unlabeled DVD was the most complete. These files were copied to P:\Surveys\Data\PacificIslands\2010_HICEAS\Oceanographic\XBT\Raw\MACII\.

Information about XBT drops, such as time, date, latitude, longitude, and comments about the profile by the at-sea oceanographer, was recorded at sea in an Excel spreadsheet, called Oceanographic_Elog_HICEAS_2010_MACII_leg#.

Perl program XBTRawFileInfo was used to get a summary of the raw files. There were three files whose drop numbers were incorrect. Those files were corrected.

Data Checks

Position Check

Perl program XBTPositionCheck compares the positions and times of the XBT drops to positions and times in the final TSG "Track Check" file. The program reads the following information from its .ini file: the location of the EDF files, the location of the final TSG Track Check file, and the number of seconds to search the TSG file for a record that matches each XBT file's time. Positions for stations without corresponding TSG data are checked against the XBT Log and ODL.

For HICEAS 2010, there were 52 drops that didn't have nearby TSG data. These were reviewed, and found to be when the TSG had not recorded any data.

Station and Drop Lists

Perl program XBTPositionAndDrops creates the Station and Drops files. From its ini file, it reads the location of the EDF files, and the location of the time offsets file. The Drop file is simply a list of the EDF Files. The Station file lists additional information for each drop: time and location, number of data points, probe type, start and end depths, and drop rate coefficients in the EDF file.

For HICEAS 2010, there were 422 drops, all using Deep Blue probes.

Profile Review and Edit

At the lab, profiles were examined by Paul Fiedler using the Visual Basic program ProfileViewer, which is located in

P:\Surveys\Data_Processing_Resources\Oceanographic\XBT\ProfileViewer. The reviewer decided what edits need to be made, and recorded that information in file XBTEdit10me.txt. Each XBT drop has a line in that file, with the EDF file name. If the profile requires any editing, the reviewer added codes for the edits to be applied. Codes are based on Bailey et al. (Quality Control Cookbook for XBT Data, CSIRO Marine Laboratories Report 221, 1994). The Quality Control Cookbook can be found in

P:\Surveys\Data_Processing_Resources\Oceanographic\XBT\ . Codes are as follows, where ddd.d is the depth exactly as recorded in the .EDF file:

- | | |
|-----------------|---|
| >RJ | Reject profile. |
| >NN | The profile is not suitable for submission to the NCEI World Ocean Database, but is good enough for deriving surface temperature and mixed layer depth, and perhaps thermocline variables, for MMTD purposes. |
| >XD ddd.d ddd.d | Exclude data within the profile. Reject data between the first and second specified depths. |

- >BD ddd.d Exclude data at the bottom end of the profile. Reject data deeper than ddd.d.
- >FT ddd.d False trigger, data recorded before the probe entered the water. This error is rare. Reject all data above ddd.d and change the depth at ddd.d to 0.7m, which is the starting depth for all drops (and then increment the depth estimates per Hanawa et al. 1995).
- >ST ddd.d Surface transient, identified by warming or cooling as the probe temperature equilibrates. Reject data before depth ddd.d. All temperatures at depths ≤ 4.0 m are rejected to routinely dismiss surface transients, so this flag is only effective if a depth >4.0 m is indicated. If the profile also contains an FT error, ddd.d is an integer that indicates the number of records after the FT correction that should be rejected.

Profile rejections and data exclusions are usually explained in a comment following an apostrophe ('): wb = wire break, ws = wire stretch, hb = the probe hit the bottom, ip = insulation penetration.

Apply Profile Edits

Perl program XBTEdit reads the editing codes entered by the reviewer, and creates edited output files. From its ini file, it reads: The location of the edit file (created in the previous step), the location of the time offsets file, the location of the raw XBT files, and the coefficients for the fall rate equation. It creates output files with the editing codes applied. (e.g. Drops marked with RJ are excluded, depths marked with XD are removed, etc.)

The program reported the following:

```

422      lines read from input file
        P:\Surveys\Data\PacificIslands\2010_HICEAS\Oceanographic\XBT\Current_Version\Inter
        mediate\MACII\XBTEdit10me.txt
0       lines started with an apostrophe, so were ignored
5       lines had an RJ code, so were ignored

417     records written to
        P:\Surveys\Data\PacificIslands\2010_HICEAS\Oceanographic\XBT\Current_Version\Inter
        mediate\MACII\XBTEdit_2010_HICEAS_MACII.dat

4       lines were not good enough for NODC, so their drop numbers were negated
413     records written to
        P:\Surveys\Data\PacificIslands\2010_HICEAS\Oceanographic\XBT\Current_Version\Inter
        mediate\MACII\XBTNODC_2010_HICEAS_MACII_6.691_0.00225.dat

11.89   Average temperature of all 477785 output points whose drops met the Min To Avg
        depth
23.76   Average temperature of all 60734 output points <= 100 m whose drops met the Min To
        Avg depth
        Note that this is based on depths calculated using the Hanawa et al 1995 formula

```

Correct Depth and Temperature

Perl program XBTCorrect adjusted XBT depths and temperatures as described in P:\Surveys\Data_Processing_Resources\Oceanographic\XBT\CH XBT fall-rate method.pdf. The program reported the following:

```
Reading XBTCorrect.ini...
Reading CH14 table 'P:\Surveys\Data_Processing_Resources\Oceanographic\XBT\Fall
rate\CH14_table1.txt'...
Reading CH14 table 'P:\Surveys\Data_Processing_Resources\Oceanographic\XBT\Fall
rate\CH14_table2.txt'...
Processing
P:\Surveys\Data\PacificIslands\2010_HICEAS\Oceanographic\XBT\Current_Version\Inter
mediate\MACII\XBTEdit_2010_HICEAS_MACII.dat...
Deep Blue: Thermal Bias = 0.02305, A = 6.640, B = 0.00248, Offset = 2.047

418   lines read from
      'P:\Surveys\Data\PacificIslands\2010_HICEAS\Oceanographic\XBT\Current_Version\Inter
mediate\MACII\XBTEdit_2010_HICEAS_MACII.dat'
0     lines started with an apostrophe, so were ignored
417   records written to
      P:\Surveys\Data\PacificIslands\2010_HICEAS\Oceanographic\XBT\Current_Version\Inter
mediate\MACII\XBTCorrect_2010_HICEAS_MACII_Deep Blue.dat

4     had negative drop numbers, which means their data was not good enough for NODC
```

Final data files

The NODC file created in the XBTEdit step was copied to \Final and was renamed as:
XBTArchive_2010_HICEAS_MACII_Deep Blue.dat

Each drop record in a XBTArchive file starts with fields under the following column headers:

```
UTC Date, UTC Time, Time Offset, Local Date, Local Time, Latitude, Longitude,
Drop #, Probe Type, # Points, H95 FRE Coefficients...
```

The column headers after that are depths (m), based on the Hanawa et al 1995 fall rate equation, e.g:

```
..., 0.7, 1.3, 2.0, 2.7, 3.3, 4.0, 4.7, 5.4, 6.0, 6.7, 7.4, 8.0, 8.7,...
```

The drop records data in those columns are temperatures (°C) at those depths. Missing data (e.g., surface transients) are blank.

The file that had depth and temperature corrected based on Cheng et al was also copied to \Final. This file is:

```
XBTCorrect_2010_HICEAS_MACII_Deep Blue.dat
```

Each drop record in a XBTCorrect file starts with fields under the following column headers:

```
UTC Date, UTC Time, Time Offset, Local Date, Local Time, Latitude, Longitude,
Drop #, Probe Type, CH14 Values, # Points...
```

The column headers after that are depths (m), based on the Cheng et al fall rate correction, e.g:

..., 0.6, 1.3, 1.9, 2.6, 3.3, 3.9, 4.6, 5.3, 5.9, 6.6, 7.2, 7.9, 8.6, 9.2, ...

Paul Fiedler
Dan Prosper
24 May 2021